

References and Literature Grading

CTA of the Brain Is a Reasonable Option to Consider to Help Rule out Subarachnoid Hemorrhage in Select Patients (12/12/2014)

1. Kowalski RG, Claassen J, Kreiter KT, et al. Initial misdiagnosis and outcome after subarachnoid hemorrhage. *JAMA*. 2004; 291:866–9. (Grade C)
2. Perry JJ, Stiell I, et al. Clinical Decision Rules to Rule Out Subarachnoid Hemorrhage for Acute Headache. *JAMA*. 2013;310(12):1248-1255. (Grade C)
3. Perry JJ, Spacek A, Forbes M, et al. Is the combination of negative computed tomography result and negative lumbar puncture result sufficient to rule out subarachnoid hemorrhage? *Ann Emerg Med*. 2008; 51:707–13. (Grade C)
4. Shah KH, Edlow JA. Distinguishing traumatic lumbar puncture from true subarachnoid hemorrhage. *J Emerg Med*. 2002; 23:67–74. (Grade E)
5. Shah KH, Richard KM, Nicholas S, Edlow JA. Incidence of traumatic lumbar puncture. *Acad Emerg Med*. 2003; 10:151–4. (Grade D)
6. Perry JJ, Stiell IG, Sivilotti ML, Bullard MJ, Emond M, Symington C, et al. Sensitivity of computed tomography performed within six hours of onset of headache for diagnosis of subarachnoid haemorrhage: prospective cohort study. *BMJ*. Jul 18 2011;343:d4277 (Grade C)
7. Morgenstern LB, Luna-Gonzales H, Huber JC Jr, et al. Worst headache and subarachnoid hemorrhage. *Ann Emerg Med*. 1998; 32:297-304. (Grade C)
8. Byyny RL, Mower WR, Shum N, Gabayan GZ, Fang S, Baraff LJ. Sensitivity of noncontrast cranial computed tomography for the emergency department diagnosis of subarachnoid hemorrhage. *Ann Emerg Med*. 2008; 51:697–703. (Grade D)
9. Boesiger BM, Shiber JR. Subarachnoid hemorrhage diagnosis by computed tomography and lumbar puncture: are fifth generation CT scanners better at identifying subarachnoid hemorrhage? *J Emerg Med*. 2005; 29:23–7. (Grade D)
10. Lourenco AP, Mayo-Smith WW, Tubbs RJ, Sidman R. Does 16-detector computed tomography improve detection of non-traumatic subarachnoid hemorrhage in the emergency department? *J Emerg Med*. 2009; 36:171–5. (Grade D)
11. Agid R, Lee SK, Willinsky RA, Farb RI, terBrugge KG. Acute subarachnoid hemorrhage: using 64-slice multidetector CT angiography to “triage” patients’ treatment. *Neuroradiology*. 2006; 48:787–94. (Grade C)
12. Robert F. McCormack, MD, and Alan Hutson, PhD , Can Computed Tomography Angiography of the Brain Replace Lumbar Puncture in the Evaluation of Acute-onset Headache After a Negative Noncontrast Cranial Computed Tomography Scan? *Academic Emergency Medicine* Volume 17, Issue 4, pages 444–451, April 2010. (Grade E)